

BGR-1 D-12 C
BGR-1 D-12 C Margulan Erlanovich Ismoldayev

**Theory
Hodograph**
Cover sheet

Please return this cover sheet together with all the related question sheets.

Hodograph (15 points).

In curvilinear motion of a planet around a star, the direction of the velocity vector changes continuously. This can be represented by a so-called "trajectory in velocity space" and is obtained as follows: for each point on the spatial trajectory, the corresponding velocity vector is drawn so that its starting point is at the origin of the velocity space, and its magnitude and direction is the same as the velocity vector at that point. The tip of this variable velocity vector generates a curve in velocity space. (The name 'hodograph' was given to this curve by Hamilton in 1846.)

As an example, see figures 1 and 2 below. For a circular orbit (Figure 1), the magnitude of the velocity is constant and therefore, the hodograph (Figure 2) of the velocity vector for Keplerian circular motion is also a circle, the center of which is located at the origin of the velocity space. The radius of this circle is equal to the constant magnitude of the circular velocity.




Fig. 1 Spatial trajectory of the Planet with Uniform Circular Motion around the star.




Fig. 2 Corresponding hodograph

- 12.1** Write an expression for the radius of the hodograph in Fig. 2, as a function of the mass M of the star, and the radius R of the circular orbit of the planet's motion. 1.0pt

- 12.2** For a planet in a Keplerian trajectory, write the expression for centripetal acceleration vector (\vec{a}) and the magnitude of angular momentum (L). For any Keplerian trajectory, it is true that 4.0pt

$$|\Delta v| = k\Delta\theta \quad (1)$$

Where k is a constant for each type of Keplerian trajectory. Find the expression for the constant k as a function of the masses M and m of the star and the planet, respectively, and the angular momentum, L .

(Eq.1) allows us to conclude that for any Keplerian trajectory, the hodograph (v as a function of θ) is a circle, but except for circular motion, the centre of the hodograph does not coincide with the star. It is not necessary to prove this result, you may simply accept it as a given. For the hodograph of uniform circular motion, the compliance with (eq.1) is completely obvious, as evidenced in Fig. 3




Fig. 3

- 12.3** Determine the expression of the constant k for the hodograph of circular planetary motion. 2.0pt

- 12.4** Given that the hodograph of the Keplerian elliptical motion is a circle, determine the radius of this hodograph and the distance between the center of the hodograph and the position of the star, as a function of the velocities at periastron and apastron. Draw a rough sketch of the hodograph in the answer sheet as per the schematic shown in Fig. 4. The black circle is the star. 4.0pt





Fig. 4

- 12.5** Similarly, for the parabolic Keplerian trajectory, determine the radius of the corresponding hodograph and the distance from the center of that hodograph circle to the star. Express the radius as a function of the velocity at periastron. Draw a rough sketch of the hodograph circle in the answer sheet. 4.0pt


Ходограф (15 точки).

При криволинейното движение на една планета около звезда, посоката на вектора на скоростта непрекъснато се променя. Това може да представено чрез така наречената "траектория в пространството на скоростите". Тя се получава по следния начин: за всяка точка от пространствената траектория се рисува съответният вектор на скоростта така, че началото му се намира в началната точка на пространството на скоростите, а големината и посоката на вектора е същата, както и на вектора на скоростта за съответната точка. Краят на вектора на скоростта описва крива в пространството на скоростите. (Названието на тази крива "ходограф" е било дадено през 1846 г. от Уилям Хамилтън.)

Като пример вижте фигурите 1 и 2 по-долу. За кръгова орбита (Фигура 1), големината на скоростта е константа и следователно ходографът (Фигура 2) на вектора на скоростта също е окръжност, центърът на която е в началната точка на пространството на скоростите. Радиусът на този кръг е равен на постоянната големина на кръговата скорост.



Фиг. 1 Пространствена траектория на планета с постоянна кръгова скорост около една звезда.



Фиг. 2. Ходографът на скоростта на планетата.


- 12.1** Напишете израз за радиуса на ходографа на Фиг.2. като функция на масата M на звездата и на радиуса R на кръговата орбита на планетата. 1.0pt

- 12.2** За планета с кеплерова орбита напишете израз за центростремителното ускорение (\vec{a}) и за големината на момента на импулса (L). За всяка кеплерова орбита е изпълнено 4.0pt

$$|\Delta v| = k\Delta\theta \quad (1)$$

Където k е константа за всеки тип кеплерова орбита. Напишете израз за константата k като функция на масата на звездата M , планетата m и на момента на импулса, L .


Уравнението (Eq.1) ни позволява да заключим, че за всяка кеплерова орбита ходографът (v as a function of θ) е окръжност, но, с изключение на случая на кръгово движение, центърът на ходографа не съвпада със звездата. Не е необходимо да доказвате този резултат. Може просто да го приемете за даденост. За случая на равномерно кръгово движение съответствието, предвид уравнение (eq.1) е съвършено очевидно, което се потвърждава от Фиг. 3.



Фиг. 3

- 12.3** Напишете израз за константата k за ходографа на движение по кръгова орбита. 2.0pt

- 12.4** Имайки предвид, че ходографът на движението по елиптична кеплерова орбита е кръг, определете радиуса на този ходограф и разстоянието между центъра на ходографа и положението на звездата като функция на скоростите в периастръ и апоастръ. Нарисувайте примерна схема на ходографа на листа за отговори съгласно схемата, показана на Фиг. 4. Черното кръгче на схемата е звездата.



Фиг. 4

- 12.5** Аналогично определете радиуса на ходографа на кеплерова параболична орбита, както и разстоянието от центъра на окръжността на ходографа до звездата. Изразете радиуса като функция на скоростта в периастръ на орбитата. Нарисувайте на листа за отговори приблизителна схема на окръжността на ходографа.

Theory



BGR-1 D-12 A-1

A12-1

English (Official)

Hodograph (15 points).

12.1 (1.0 pt)

12.2 (4.0 pt)

$\vec{a} =$

$L =$

$k =$

12.3 (2.0 pt)

$k =$

12.4 (4.0 pt)

DELEGATION PRINT

Theory



BGR-1 D-12 A-2

A12-2
English (Official)

12.5 (4.0 pt)

DELEGATION PRINT

Theory



BGR-1 D-12 W-1

W12-1

DELEGATION PRINT

Theory



BGR-1 D-12 W-2

W12-2

DELEGATION PRINT

Theory



BGR-1 D-12 W-3

W12-3

DELEGATION PRINT

Theory



BGR-1 D-12 W-4

W12-4

DELEGATION PRINT